

My physician recommends...

# Children's Advil® Relief you can trust

The #1 selling children's fever brand\*



Children's Advil®

Lasts Up To **8** Hours

Measure dosing only with the device that comes with the product



Available in 4 flavors!  
 • Dye-Free  
 • White Grape  
 • Fruit  
 • Grape  
 • Blue Raspberry



Based on weight and age, your healthcare professional has indicated below which Advil® medicine is right for your child.

		Infants' Advil® Concentrated Drops for ages 6-23 months	Children's Advil® Suspension for ages 2-11 years	Junior Strength Advil® Chewables for ages 6-11 years	Junior Strength Advil® Tablets for ages 6-11 years
<b>Active ingredient</b>		Ibuprofen 50 mg (NSAID) <sup>†</sup> in each 1.25 mL	Ibuprofen 100 mg (NSAID) <sup>†</sup> in each 5 mL	Ibuprofen 100 mg (NSAID) <sup>†</sup> in each tablet	Ibuprofen 100 mg (NSAID) <sup>†</sup> in each tablet
<b>Indications</b>		Fever reducer; relieves minor aches and pains due to the common cold, flu, headaches, and toothaches	Fever reducer; relieves minor aches and pains due to the common cold, flu, sore throat, headaches, and toothaches	Fever reducer; relieves minor aches and pains due to the common cold, flu, sore throat, headaches, and toothaches	Fever reducer; relieves minor aches and pains due to the common cold, flu, sore throat, headaches, and toothaches
<b>Weight</b>					
<b>Age</b>					
Patient Dosing	under 6 months	<b>Do not use</b>	<b>Do not use</b>	<b>Do not use</b>	<b>Do not use</b>
	12-17 pounds	6-11 months <b>1.25 mL<sup>‡</sup></b> every 6-8 hrs not to exceed 4 doses/24 hrs	Use Infants' Advil® Concentrated Drops	Use Infants' Advil® Concentrated Drops	Use Infants' Advil® Concentrated Drops
	18-23 pounds	12-23 months <b>1.875 mL<sup>‡</sup></b> every 6-8 hrs not to exceed 4 doses/24 hrs	↓	↓	↓
	24-35 pounds	2-3 years Use Children's Advil® Suspension	<b>1 tsp<sup>‡</sup></b> every 6-8 hrs not to exceed 4 doses/24 hrs	Use Children's Advil® Suspension	Use Children's Advil® Suspension
	36-47 pounds	4-5 years ↓	<b>1½ tsp<sup>‡</sup></b> every 6-8 hrs not to exceed 4 doses/24 hrs	↓	↓
	48-59 pounds	6-8 years Use Children's Advil® Suspension, or Junior Strength Advil® Chewables or Tablets	<b>2 tsp<sup>‡</sup></b> every 6-8 hrs not to exceed 4 doses/24 hrs	<b>2 tablets</b> every 6-8 hrs not to exceed 4 doses/24 hrs	<b>2 tablets</b> every 6-8 hrs not to exceed 4 doses/24 hrs
	60-71 pounds	9-10 years ↓	<b>2½ tsp<sup>‡</sup></b> every 6-8 hrs not to exceed 4 doses/24 hrs	<b>2½ tablets</b> every 6-8 hrs not to exceed 4 doses/24 hrs	<b>2 tablets</b> every 6-8 hrs not to exceed 4 doses/24 hrs
	72-95 pounds	11 years ↓	<b>3 tsp<sup>‡</sup></b> every 6-8 hrs not to exceed 4 doses/24 hrs	<b>3 tablets</b> every 6-8 hrs not to exceed 4 doses/24 hrs	<b>3 tablets</b> every 6-8 hrs not to exceed 4 doses/24 hrs
	12+ years	Use Advil® Tablets or Liqui-Gels®	Use Advil® Tablets or Liqui-Gels®	Use Advil® Tablets or Liqui-Gels®	Use Advil® Tablets or Liqui-Gels®

Use the fever diary below to record your child's fevers and dosing schedule.

Fever: _____ °F or _____ °C	Time: _____	Dose: _____
Fever: _____ °F or _____ °C	Time: _____	Dose: _____

<sup>‡</sup> 1 Tsp=100 mg ibuprofen per 5 mL.  
 \*Among over-the-counter brands based on 52 weeks ending 9/4/11.  
<sup>†</sup>Nonsteroidal anti-inflammatory drug.  
<sup>‡</sup>Measure dosing only with the device provided.

Please refer to product label for complete information and warnings.